

A következő statisztikákat a

**Practice Guidelines for the Treatment
of Unwanted Same-Sex Attractions and Behavior¹**

National Association for Research and Therapy of Homosexuality,
Task Force on Practice Guidelines for the Treatment of
Unwanted Same-Sex Attractions and Behavior²
Salt Lake City, Utah

c. könyvből vettük ki.




Figure 1. Mood Disorders, OR 2.78 SSA men, 1.48 SSA women (Tjepkema, 2008; Canada). Bisexual for male and female from the same reference is also given.

Anxiety Disorder




Figure 2. Anxiety Disorder, OR 2.83 men, 1.50 women (Tjepkema, 2008; Canada).

General Anxiety Disorder




Figure 3. Generalized Anxiety Disorder, OR 2.8 men, 1.2–6.5 women. The figures for women are a range for a variety of diagnostics; in this paper means will be plotted (Fergusson, Horwood & Beautrais, 1999; New Zealand longitudinal study).

Depression




Figure 4. Depression, OR 3.5 (Cochran, Sullivan, & Mays, 2003; United States).

Simple Phobia




Figure 5. Simple Phobia, OR 3.61 men, 1.27 (NS) women; latter not plotted (Cochran et al., 2003).

Promiscuity




Figure 6. Promiscuity, OR 2.75 men, 4 women (Laumann, Gagnon, Michael, & Michaels, 1994; United States). Medians are used. The issue is whether hypersexuality exists; a later discussion on sexual compulsivity indicates this is likely.

Partner Interpersonal Violence




Figure 7. Violence in same- and opposite-sex couples, OR 3.0 men, 1.5 women